

Turmeric Extract As An Antioxidant In Repeatedly Used Cooking Oil

Andi Abriana, Eva Johannes

ABSTRACT: One of the natural antioxidant sources is turmeric plant from zingiberaceae family that contains curcumin. Curcumin is an important component isolated from turmeric (*Curcuma longa*), that is used as traditional medicines and food additive substance. This study was aimed to examine the use of turmeric as an antioxidant in repeatedly used cooking oil and to find out the mechanism by which the turmeric extract inhibits the formation of trans fatty acid in repeatedly used cooking oil. This study was an experimental study in which the turmeric extract was obtained by maceration. The obtained turmeric extract was then added into cooking oil that will be used later to fry repeatedly for 10 times. Frying treatments consisted of frying without turmeric extract addition (control) and frying with turmeric extract addition at concentrations of 0.02%, 0.03%, and 0.04%. The samples of the used cooked oil were collected at frying 1, frying 5, and frying 10 and analyzed for their trans fatty acid content by gas chromatography. Study findings indicate that turmeric extract addition into cooking oil reduced the formation of trans fatty acid and extended the use time of cooking oil. According to trans fatty acid content analysis, the best treatment was the addition of turmeric extract at concentration of 0.03%.

Keywords: Turmeric extract, antioxidant, cooking oil, repeated frying.

INTRODUCTION

Fried foods are the kind of foods that contains fat and cholesterol that frequently induce the development of various diseases such as coronary diseases and others. Consuming fried foods commonly sold in streets and many places is risky. Fried foods are commonly fried with repeatedly used cooking oil at high temperature and in long time (deep frying). Chemically, cooking oil that has been used repeatedly is very different from fresh palm oil that has not been used for frying. In palm oil there are about 45,5% of saturated fatty acid dominated by palmitatefatty acid and about 54.1% of unsaturated fatty acid dominated by oleic fatty acid. Whereas in palm oil that has been used repeatedly the fatty acid (trans fatty acid) is much more higher than the saturated fatty acid. Trans fatty acid is very harmful to health because it can induce various diseases such as cardiac diseases and stroke [1]. Some of the fried foods that contain trans fatty acid are fried bananas, fried cassavas, croquette, fried fermented soybeans, and powdered fried chickens that are fried with repeatedly used cooking oil. It is these types of food that contribute to high trans fatty acid intake. The formation of trans fatty acid by partial hydrogenation in vegetable oil that contain unsaturated fatty acid if through heating process with high temperature. Frying using heated fat or oil at high temperature of 150°C – 190°C is one of food serving technique that becomes popular because it can provide better color, taste, and texture for fries. Two methods of frying are practiced, namely (1) Shallow frying and (2) Deep frying. Shallow frying uses pan (pan frying) and the fried foods are only partly submerged in the oil.

In deep frying technique, the fried foods are submerged completely in the oil and the oil temperature can reach 200-205°C, that it needs oil in high number. Deep frying technique causes changes in stability and quality, taste, color, and texture of the fried foods and their nutrient contents. Commercially, this process is widely applied particularly for industrial scale production of various products such as fried potatoes, seafood, egg rolls, and chicken patties. Frying process with deep frying technique enables heat transfer during the process from the hot oil into cool foods. It is this feature that makes this process fast. The fried foods have delicious taste, layered by crispy surface, interested color, and the oil absorption by the fried product will result in the desired mouthfeel, easily reconstructed, and free from harmful microorganisms [2,3,4,5]. However, it is associated to health risk because the repeatedly used cooking oil will subject to oxidation that produce peroxide groups and cyclic monomer. According to [1], in addition to harmful effect by peroxide group to health, the use of repeatedly used cooking oil also produces aldehyde-type acrolein that produce itchiness in throat. Curcumin is an antioxidant with concentrated yellow color isolated from turmeric, known effective as food composition because beside it slows down the food degradation, it is also effective in inhibiting the formation of lipid peroxide compared to synthetic antioxidant such as BHT [6]. The presence of antioxidant in fat will reduce the oxidation process rate, by scavenging the free radicals and change them into non-reactive "second hand" radicals that are not harmful to health [7]. This study was conducted to determine the effect of turmeric extract as natural oxidant on the extension of use time of repeatedly used cooking oil.

METHOD

This study was conducted from March 2014 to September 2014, using experimental method with the following steps:

Turmeric extraction: 1 kg of turmeric was selected and cleansed initially. After that the turmeric were peeled and chopped and then dried under sun light for 1 week. After dried, the turmeric were crushed by using blender, weighted, and 500 g of fine turmeric powder were stored in closed container. Turmeric were extracted by maceration using ethanol (1:1) for 24 hours, and then filtered by

- Andi Abriana, Eva Johannes
- Department of Agricultural Technology, Agricultural Faculty, Bosowa 45 University Makassar, Email: andi_abriana@yahoo.com
- Department of Biology, Faculty of Mathematic and Natural Sciences, Email: evajohannes@ymail.com

buchner filter to separate the residue and its supernatant. Maceration was performed three times. The result of the maceration, in the form of supernatant, was dried by using evaporator in order that the ethanol liquid was separated from its macerate. 50 mg of thick macerated was obtained.

Food frying: The food materials used in this study were bananas (saba banana) and cooking oil. Frying was performed without the addition of turmeric extract (control) and with the addition of turmeric extract at concentrations of 0.02%; 0.03%, and 0.04%. Frying process was started by adding 3.5 liters of fresh cooking oil into frying pan, the pan was heated up to 180°C. The food materials were fried for 10 minutes and it was attempted to not frequently stir the fries to reduce convection flow from oil and oxidation reaction due to aeration process. Frying was performed up to 10 times. The samples of cooking oil were collected in frying 1, 5, and 10 (each 100 mL) to analyze the trans fatty acid content.

Trans Fatty Acid Measurement: Identification of trans fatty acid composition was performed in cooking oil that had been used to fry bananas with the samples were collected at frying 1, 5, and 10. Analysis of trans fatty acid was performed by gas chromatography (GC) to separate its forms and trans fatty acid configurations. Components were separated by evaporation, carried by inert gas and passed through a column or stationary phase in the form of solid or liquid substances that is not easy to evaporate that attached to inert supporting substance with the following ways:

- Sample preparation (hydrolysis and esterification). First, oil sample was weighed in a teflon closed tube, and added with 1 mL of 0.5 N NaOH in methanol and heated with water bath for 20 minutes.
- 2 mL of BF₃ 16 % and 5 mg/ml of internal standard were added and heated again for other 20 minutes. After cooling down, 2 mL of saturated NaCl and 1 mL of hexane was added. Hexane layer was separated and added into a tube containing 0.1 g of anhydrous Na₂SO₄ and kept for 15 minutes. Liquid phase was separated in injected into gas chromatography.
- Analysis of fatty acid components, as FAME by gas chromatography, cyanoprimethylsil column (capillary column). The instrument conditions were adjusted as following: column dimension (length= 60 m; Å depth = 0,25 mm, 0,25 Å Film Thickness); flow rate of N₂ : 20 mL/min; lajuair H₂:30 mL/min; flow rate of air: 200 Å 250 mL/min; injector temperature: 200°C; detector temperature: 230°C; column temperature: temperature program (column temperature: initial 190°C, stationary 15 min, end 230°C stationary 20 min and rate 10°C/min; ratio = 1:8; inject volume: 1 Å L; linear velocity: 20 cm/sec.
- Analysis was started from solvent injection (1 µL) into column to obtain baseline values, and then continued by injecting 5 µL of FAME standard solution. When all peaks had been out, sample was injected in the volume of 5 µL. Retention time and sample peak were measured for each component and compared to standard and calculated with the following formula:

$$C_x = \frac{A_x \cdot R \cdot C_s}{A_s}$$

Note:

- C_x : Concentration of component x
 C_s : Concentration internal standard
 A_x : Area of component x peak
 A_s : Area of internal standar peak
 R : Response

Data Analysis: Data were analyzed by using SPSS version 17 with contrast further test.

RESULTS AND DISCUSSION

Frying Without Turmeric Extract Addition (Control)

The frying of food materials with cooking oil without the addition of turmeric extract (control). Study findings indicate that the more frequent the use, the higher the trans fatty acid content in repeatedly used cooking oil. (Figure 1). According to [8], unsaturated fatty acid level is increasing with the use frequency of the repeatedly used cooking oil.

Frying with Turmeric Extract Addition

The frying of foods with addition of turmeric extract at the concentrations of 0.02%, 0.03%, and 0.04% in the used cooking oil indicated that in frying 10 there was a decrease in trans fatty acid content. This means that the addition of turmeric extract in cooking oil reduces the formation of trans fatty acid. Analysis of trans fatty acid content in the used cooking oil without addition of turmeric extract (control) indicated higher trans fatty acid content at frying 10; whereas the used cooking oil with addition of turmeric extract at concentration of 0.02%, 0.03% and 0.04% indicated reduction trans fatty acid content at frying 10 (Figure 1). This was related to curcumin that inhibit the auto-oxidation rate by modifying the lipid radical into more stable form, that commonly inhibit the fat oxidation reaction [9]. According to [10], turmeric extract is an antioxidant with good reactivity and has the potential to be used as natural antioxidant in food industry. According to [11], turmeric extract added into cooking oil can reduce trans fatty acid content, because the turmeric extract is capable of inhibiting oxidation reaction in the oil that the trans fatty acid formation is slowed down significantly (P<0.05). The presence of antioxidant added into the repeatedly used cooking oil can decelerate the oil oxidation rate during frying and contribute to sensoric acceptance of fried foods. The findings of this study are supported by the statement of [12] that the addition of antioxidant in cooking oil determine the stability of oxidation during frying; and according to Tuba and [13], curcumin is effective to be used as antioxidant because it can scavenge free radical by donating H atom from phenolic as its active group. According to [11], the use of antioxidant initially reduces the oxidative stability loss during frying, and the reduces heat degradation under the commonly used frying conditions.




Figure 1. The results of trans fatty acid analysis in cooking oil that was used to fry bananas

When compared to analysis results of trans fatty acid content with addition of turmeric extract at concentrations of 0.02%, 0.03%, and 0.04%, it can be seen that the higher the concentration of the added turmeric extract, the lower

the trans fatty acid content. The addition of turmeric extract at concentrations of 0.03% and 0.04% were not significantly different and according to organoleptic test against cooking oil and fried food colors with addition of 0.03% and 0.04% turmeric extract, it was shown that the addition of 0.04% turmeric extract resulted in very yellow color of cooking oil and the fried foods; whereas the addition of 0.03% turmeric extract resulted in less yellow color of cooking oil and the color of the fried foods is not significantly different from normal color of the fried foods. According to follow-up test by contrast test (Table 1), it was shown that the control treatment and turmeric extract addition treatment at all concentrations were different significantly. This indicates that turmeric extract can be used as an antioxidant in repeatedly used cooking oil and can inhibit the formation of trans fatty acid in repeated frying and extend the use time of cooking oil. This is in accordance to [13] that the presence of antioxidants in cooking oil is very important in stabilizing the complex frying system and improve the appearance and lifetime of fried foods. The best treatment for repeatedly used cooking oil is the addition of turmeric extract at concentration of 0.03%.

Table 1. The results of follow-up test for trans fatty acid content in used banana cooking oil

| Frying repetition | Control | Addition of 0.02% turmeric extract (% b/b) | Addition of 0.03% turmeric extract (% b/b) | Addition of 0.04% turmeric extract (% b/b) |
|-------------------|---------------------|--|--|--|
| Frying 1 | 0.103 ^{ns} | 0.11 ^{**} | 0.093 ^{**} | 0.093 ^{**} |
| Frying 5 | 0.107 ^{ns} | 0.11 ^{**} | 0.093 ^{**} | 0.093 ^{**} |
| Frying 10 | 0.12 ^{ns} | 0.09 ^{**} | 0.09 ^{**} | 0.087 ^{**} |

^{ns}) not significantly different ^{**})very significantly different

CONCLUSION

The addition of turmeric extract at the concentrations of 0.02%, 0.03%, and 0.04% reduces the formation of trans fatty acid in repeatedly used cooking oil, and extends the use time of cooking oil. The best result was obtained with addition of 0.03% turmeric extract.

Acknowledgement

The author thanks to Directorate General of Higher Education, for the financial support for this study.

BIBLIOGRAPHY

- [1] Suci D. Perwitasari., 2013. <http://www.e-jurnal.com/2013/10/penambahan-kunyit-sebagai-antioksidan.html>
- [2] Blumenthal, M.M. 1996. Frying Technology. : Bailey's Industrial Oil and Fat Technology; Edible Oil and Fat Product: Product and Application Technology (4th ed., Vol 3). Wiley-Interscience Publication. New York. pp.429-482.
- [3] Krishnamurthy, R.G. dan Vernon C. W. 1996. Salad Oil and Oil-Based Dressings.: Bailey's Industrial Oil and Fat Technology; Edible Oil and Fat Product: Product and Application Technology (4th ed., Vol 3). Wiley-Interscience Publication. New York. pp. 193-224.
- [4] Choe, E and D.B. Min. 2007. Chemistry of Deep-Fat Frying Oils. Journal of Food Science. Vol. 72. Nr. 5. Institute of Food Technologists.
- [5] Jana. Pinto Sunetaa, Patel Komal and Chaudary Apurva. 2011. Choosing The Vegetable Oil For Frying of Food – A Review. Journal of Advances in Developmental Research Vol. 2 No. 2:213-220
- [6] Suhana., 2004. Studi Isolasi dan Aktivitas Radical scavenger (penangkap radikal) Rimpang Kunyit (curcuma longa). Karya Utama Sarjana Kimia F. MIPA UI. Depok
- [7] Widjaya, C.H., 2003. Antioxidant Role on Body Health. Edisi 1V Grasinda. Jakarta.
- [8] Songklanakarin., 2004. The Use of Stable Free Radical Diphenyl picrylhydrazyl (DPPH) for estimating Antioxidant Activity. J. Scie. Tech. 26. 211-219.

- [9] Nor, F. M., Suhaila Mohamed, Nor Aini Idris and Razali Ismail. 2009. Antioxidative Properties of Curcuma longa Leaf Extract in Accelerated Oxidation and Deep Frying Studies. Journal American Oil Society (86) 141-147. Springer AOCS.
- [10] Jaswir, I., Yaakob B.Che Man and David D. Kitts. 2000. Use of Natural Antioxidants in Refined Palm Olein During Repeated Deep-Fat Frying. Food Research International Journal (33): 501-508. Elsevier Science Ltd.
- [11] Man, Yaakob B. C., Wanna A, Russly A.R and Salmah Yusof. 2003. Quality Characteristics of Refined, Bleached and Deodorized Palm Olein and Banana Chips after Deep-Fat Frying. Journal of The Science of Food and Agriculture. Society of Chemical Industry.
- [12] Tuba Ak and Ilhami Gulcin. 2008. Antioxidant and Radical Scavenging Properties of Curcumin. Chemico –Biological Interactions Journal (174) 27-37. Elsevier Ireland Ltd
- [13] Casal, S., Ricardo M, Artur S, Beatriz P.P. O and Jose A. P. 2010. Olive Oil Stability Under Deep-Frying Conditions. Food and Chemical Toxicology Journal (48):2972-2979.